



产品详情

## L-8R $\beta$ Rabbit Polyclonal Antibody

产品货号	产品名称	储存条件	保质期
IM30005	IL-8R $\beta$ Rabbit Polyclonal Antibody	-20℃	1 年

### Product information

别名	CXCR2; IL8RB; C-X-C chemokine receptor type 2; CXC-R2; CXCR-2; CDw128b; GRO/MG SA receptor; High affinity interleukin-8 receptor B; IL-8R B; IL-8 receptor type 2; CD182
产品名称	IL-8R $\beta$ Rabbit Polyclonal Antibody
类别	抗原抗体
基因名称	CXCR2
蛋白名称	C-X-C chemokine receptor type 2
推荐应用	WB, IHC-P, IF-P, IF-F, IF-ICC, ELISA
反应种属	Human, Rat, Mouse
浓度	1mg/ml
存储缓冲液	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type p reservative N.
免疫原	The antiserum was produced against synthesized peptide derived from the N-terminal region of human CXCR2. AA range: 1-50
特异性	IL-8R $\beta$ Polyclonal Antibody detects endogenous levels of IL-8R $\beta$ protein.
稀释度	WB1:500-1:2000, IHC-P1:100-1:300, ELISA1:20000, IF-P/IF-F/IF-ICC1:50-200

预测分子量	40kDa
运输及保存条件	-20° C/1 year
Contains	no preservative
Storage conditions	-20° C
宿主	Rabbit
同种型	Rabbit, IgG
背景介绍	<p>C-X-C motif chemokine receptor 2(CXCR2) Homo sapiens The protein encoded by this gene is a member of the G-protein-coupled receptor family. This protein is a receptor for interleukin 8 (IL8). It binds to IL8 with high affinity, and transduces the signal through a G-protein activated second messenger system. This receptor also binds to chemokine (C-X-C motif) ligand 1 (CXCL1/MGSA), a protein with melanoma growth stimulating activity, and has been shown to be a major component required for serum-dependent melanoma cell growth. This receptor mediates neutrophil migration to sites of inflammation. The angiogenic effects of IL8 in intestinal microvascular endothelial cells are found to be mediated by this receptor. Knockout studies in mice suggested that this receptor controls the positioning of oligodendrocyte precursors in developing spinal cord by arresting their migration. This gene, IL8RA, a gene encoding another high affinity IL8 receptor, as.</p>
组织表达	Brain, Placenta.
细胞定位	Cell membrane; Multi-pass membrane protein.

信号通路	Cytokine-cytokinereceptorinteraction;Chemokine;Endocytosis;Epithelial cell signaling in Helicobacter pylori infection;function:Receptor for interleukin-8 which is a powerful neutrophil chemotactic factor.Binding of IL-8 to the receptor causes activation of neutrophils.This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. Binds to IL-8 with high affinity.
功能	Also binds with high affinity to CXCL3, GRO/MGSA and NAP-2.,online information:CXC chemokine receptors entry,PTM:Phosphorylated upon ligandbinding;whichisrequired for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.
纯化	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal

免责声明：所有产品仅用作科学研究，不得用于其他用途！本公司将不为任何不正常使用此产品时所发生的意外负责。

北京伊事达科技有限公司

电话：13564444959

官网：[www.followme-shop.com](http://www.followme-shop.com)

地址：北京市海淀区东北旺西路58号尚科办公社区C区一楼



公众号



客服