

Tubulin beta-3(9117) Mouse Monoclonal Antibody



产品货号	产品名称	储存条件	保质期
IM12060	Tubulin beta-3(9117) Mouse Monoclonal Antibody	-15℃to-25℃	1年

产品详情:

产品货号	IM12060	
别名	TUBB3;TUBB4;Tubulin beta-3 chain;Tubulin beta-4 chain;Tubulin	
	beta-III	
产品名称	Tubulin beta-3(9117)Mouse Monoclonal Antibody	
类别	抗体产品	
基因名称	TUBB3	
蛋白名称	Tubulin beta-3 chain	
推荐应用	WB, IF-ICC, ELISA	
反应种属	Human, Mouse, Rat	
存储缓冲液	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA	
Human Gene ID	10381	
Human Swissprot	Q13509	
INO.		
Mouse Gene ID	22152	
Mouse Swissprot	Q9ERD7	
No.		



Rat Gene ID	246118	
Rat Swissprot	Q4QRB4	
免疫原	Synthesized peptide derived from human β III Tubulin AA range:400-451	
特异性	This antibody detects endogenous levels of Tubulin β3.	
稀释度	WB 1:1000-5000. IF-ICC 1:100-500. ELISA 1:5000-50000	
参考分子量	55kDa	
预测分子量	50kDa	
运输及保存条件	-15°Cto-25°C/1 year(Do not lower than -25°C)	
宿主	Mouse, Monoclonal/IgG2b, kappa	
背景介绍	tubulin beta 3 class III (TUBB3) Homo sapiens This gene encodes a class III member of the beta tubulin protein family. Beta tubulins are one of two core protein families (alpha and beta tubulins) that heterodimerize and assemble to form microtubules. This protein is primarily expressed in neurons and may be involved in neurogenesis and axon guidance and maintenance. Mutations in this gene are the cause of congenital fibrosis of the extraocular muscles type3. Alternate splicing results in multiple transcript variants. A pseudogene of this gene is found on chromosome 6. [provided by RefSeq, Oct 2010]	
细胞定位	Cytoplasmic	
表达	Expression is primarily restricted to central and peripheral nervous system. Greatly increased expression in most cancerous tissues.	



	domain: The highly acidic C-terminal region may bind cations such as	
	calcium.function:Receptor for MSH(alpha, beta and gamma) and ACTH.	
	The activity of this receptor is mediated by G proteins which activate	
	adenylate cyclase.function:Tubulin is the major constituent of	
	microtubules. It binds two moles of GTP, one at an exchangeable site	
	on the beta chain and one at a non-exchangeable site on the	
功能	alpha-chain.polymorphism:Genetic variations in MC1R are associated	
	with variation in skin/hair/eye pigmentation type	
	2(SHEP2)[MIM:266300]. Hair, eye and skin pigmentation are among the	
	most visible examples of human phenotypic variation, with a broad	
	normal range that is subject to substantial geographic	
	stratification. In the case of skin, individuals tend to have lighter	
	pigmentation with increasing distance from the equator.	
纯化	Protein G	

注意事项:

- 1. 本产品仅供科研使用。请勿用于医药、临床诊断或治疗,食品及化妆品等用途。请勿存放于普通住宅区。
- 2. 为了您的安全和健康,请穿好实验服并佩戴一次性手套和口罩操作。
- 3. 实验结果可由多种因素影响,相关处理只限于产品本身,不涉及其他赔偿。

免责声明: 所有产品仅用作科学研究,不得用于其他用途! 本公司将不为任何不正常使用此产品时所发生的意外负责。

北京伊事达科技有限公司

电话: 13564444959

官网: www.followme-shop.com

地址:北京市海淀区东北旺西路58号尚科办公社区C区一楼





公众号

客 服